Evaluation of Oral and Dental Hygiene in Students: Film- and Lecture-based Educational Intervention

Reza Jorvand¹, Omidali Gholami²*, Fatemeh Khirolahi³, Hojat Hasani⁴, Kobra Momeni⁵, Morteza Mansourian⁶

Abstract

Aim: Oral health is among the important components of individual general health that affects the health of entire body. This study aims at investigating the effect of educational intervention on the primary school students' oral health.

Methods: This quasi-experimental study was conducted on 175 primary school students recruited randomly from different regions of Dehloran City during the school year of 2014-15. The testimonial was obtained after holding briefings for the parents. Data were collected using a researcher-made questionnaire, which was administrated at baseline and 3 months after the intervention. The content validity was approved by the expert panel and its reliability was confirmed by Cronbach's alpha coefficient. Data were analyzed by SPSS software 18, inferential statistical tests, and paired t-test. This study was supported by the HSR Council and the Ethics Committee of Ilam University of Medical Sciences.

Findings: The results indicated significant changes in the samples' knowledge (from 3.6 ± 1.37 to 4.98 ± 1.57), attitude (from 28.17 ± 5.7 to 30.98 ± 6.35) and behavior (15.03 ± 2.98 to 17.08 ± 2.60) after the intervention.

Conclusion: The positive results of educational intervention indicate that the education through appropriate methods can improve students' behavior in the field of oral health. Therefore, it is suggested more emphasis on oral health education at school age.

Keywords: Educational intervention, Dental hygiene, Students

Email: kobramomeni@ut.ac.ir

^{1.} Ph.D. Candidate, Department of Health Education and Health Promotion, Faculty of Health, Ilam University of Medical Sciences, Ilam, Iran Email: rezajorvand@yahoo.com

^{2.} M.Sc., Department of Public Health, Center for Health, DEHLORAN City, Ilam University of Medical Sciences, Ilam, Iran Email: Omidgholami88@yahoo.com

^{3.} Oral Health Expert, Department of Public Health, Center for Health, DEHLORAN City, Ilam University of Medical Sciences, Ilam, Iran Email: kheirollaheefatemeh@gmail.com

^{4.} School Health Expert, Department of Education, DEHLORAN, Ilam Province, Iran

Email: Hassanihojat@ yahoo.com

^{5.} M.Sc. Student, Department of Biostatistics, Tehran University, Tehran, Iran

^{6.} Assistant Professor, Department of Health Education and Health Promotion, Faculty of Health, Iran University of Medical Sciences, Tehran, Iran Email: mansourian55@gmail.com

Introduction

Mouth is a mirror of the body, and teeth are the main components [1]. The oral cavity has many functions associated with the daily living of people such as eating, beauty, speech and social interactions [2]. Oral diseases such as tooth decay and periodontal are among the most common diseases worldwide [3-5], and poor oral and teeth hygiene not only causes diseases related to the oral cavity but also can be a threat to the vital body systems and even lead to death [6]. World Health Organization (WHO) has focused on the use of experiences and evaluation of previous objectives with an emphasis on the importance of oral health as an indispensable component of public health until 2020 [7].

Tooth decay has been climbing in our country; this is likely due to poor public awareness of the role and importance of oral hygiene in maintaining dental health [8]. In order to reduce public oral health problems and mitigate the costs of subsequent treatment, additional efforts should be made in order to raise the awareness, attitude, and enhanced performance of people, especially children, who are in the course of learning, and intellectual and cultural growth and development [9]. Oral health education at schools as a small but important social institution with diverse training needs and systematic planning tailored to each country is of fundamental interest to organizations

providing health, as well as a recommendation of WHO. Hence, one of the principles in PHC, The Alma Ata Declaration, has been health promotion through education to reduce unnecessary health inequalities considered in the Ottawa conference [10].

In domestic literature review, few studies were found in which the impact of using video tutorials on the oral health of students was investigated [10-12]. However, the dramatic ways in addition to the educational aspect are used to create or modify attitudes and skill training in health education programs [13].

Therefore, due to the high rate of tooth decay at schools [14], the present study was implemented to investigate the effect of combined educational intervention including lecture and film screening on the knowledge, attitude, and behavior of elementary students about oral health in Dehloran City during the academic year 2014-15.

Objective

Oral health is among the important components of individual general health, and affects the health of the entire body. This study aims at investigating the effect of educational intervention on the primary school students' oral health.

Materials and Methods

This study was performed on 175 students of

fourth and fifth grades at an elementary school in Dehloran City during the academic year 2014-15. The samples were selected through random cluster sampling (in two schools and six classrooms). The inclusion criteria were studying at the fourth or fifth grade of elementary schools in Dehloran City, filling in a consent form by the parents, participating in the training courses, and cooperating until the end of the project. The drop-out students and those who did not participate in the educational programs were excluded from the study.

The data collection tool was a self-made questionnaire with 39 questions including 4 parts of the personal information (9 questions), knowledge (10 questions by scoring zero and one), attitude (10 questions or 5-point Likert scale and scoring 1-5), and behavior (10 questions by scoring 0-4).

The initial questionnaire was developed according to the research purposes based on credible literature. The questionnaire's validity was assessed through the content validity by the comments of experts in health and dentistry education. Cronbach's alpha determined the reliability of the questionnaire with a score of 0.75. Upon coordinating with the authorities, the students' parents were invited to attend a briefing, and complete the consent forms. The first phase questionnaires were collected before the educational intervention. The questionnaires were completed by trained

dental hygienists through interview method. The contents of intervention were provided based on the results of the first stage by relying on the initial assessment of requirements and literature.

Twenty five elementary students participated in the pilot study. In this study, the students' opinions about the best way to learn were collected in addition to investigating the reliability of the tools. Based on the results of the pilot study, film screening and training was selected with the help of experts as the transmission method of educational content.

The educational intervention included screenings (from production of the Oral Hygiene and Health Unit of the Health Center of Ilam Province) for students and providing additional details by the dentist. The training was conducted three times during a month.

The video composed of an introduction to the oral parts, layers and surfaces of the mouth and teeth, showing healthy and unhealthy gums, the use of proper toothbrush and toothpaste, the right way of brushing and flossing and the use of sodium fluoride mouthwash, as well as additional details about the side effects of not observing oral hygiene, regular visit to the dentist, healthy nutrition and so on presented by the dentist. Three months after the intervention, the questionnaires were completed by referring back to the respective schools. Data were analyzed using the statistical software SPSS 18 by independent sample t-test and correlation coefficients. Each student was donated a toothbrush and a toothpaste to ensure they have access to them. In addition, this study was approved by the HSR Council and the Medical Ethics Committee of Ilam University of Medical Sciences.

Results

Girls represented 48% of our subjects. The

mean age of the samples was 10.55±0.78 years. Most of the students (53.7%) were in the fifth grade, and the average of the students' family size was 3.71. Majority of the mothers were housewives (70.9%), while only 33.7% were employees. While the education (26.9 percent) of the fathers was at the bachelor's level, and at the middle school's level for 30.9% of the mothers. The usage rate of toothbrush and dental floss was 59% and 36%, respectively (Table 1).

Table 1: Distribution of the demographic characteristics of study participants

Variable		Frequency (N = 175)	Percent
Condon	Girl	84	48
Genuer	Boy	91	52
Age	M ean \pm SD	10.55 ± 0.778	
Grade	Fourth	81	46.3
	Fifth	94	53.7
Children sequence	First	81	22.3
	Second	39	17.1
	Third	30	9.7
	Fourth	17	9.7
	Fifth and more	8	4.6
Number of family members	Two	1	0.6
	Three	8	4.6
	Four	70	40
	Five	55	31.4
	Six and more	41	23.4
Mother's job	Housewife	124	70.8
	Employee	11	6.3
	Teacher	29	16.6
	Self-employed	11	6.3
Father's job	Teacher	20	11.4
	Employee	59	33.7
	Labor	25	14.3
	Self-employed	57	32.6
	Unemployed	14	8
Mother's education	Illiterate	28	16
	Guidance	54	30.9
	Diploma	43	24.6
	Associate degree	20	11.4
	Bachelor's	30	17.1
Father's education	Illiterate	18	10.3
	Guidance	38	21.7
	Diploma	45	25.7
	Associate degree	27	15.4
	Bachelor's	47	26.9

Moreover, the mean scores of knowledge, attitude, and behavior of the students changed

significantly following the intervention (Table 2).

l I	ariable	Mean (SD)	df	Sig. (2-tailed)	
Knowledge	Before intervention	3.60 ± 1.37	1/10	P = 0.000	
Kilowieuge	After intervention	4.98 ± 1.57	14)	1 = 0.000	
Attitude	Before intervention	28.16 ± 5.7	173	P = 0.000	
	After intervention	32.18 ± 8.24	175		
Behavior	Before intervention	15.03 ± 2.98	145	D = 0.000	
	After intervention	17.08 ± 2.60	143	r = 0.000	

Table 2: Mean scores of knowledge, attitude and behavior of the participants before and after intervention

Statistical tests showed no statistically significant relationship between gender, age and number of children in the family of participants and the students' knowledge, attitude and behavior. While studying the relationship between the classroom of participants and the study variables, it was found that there was a significant relationship just between classroom and behavior. In the present study, no significant relationship was found between the education level of parents and the knowledge, attitude and behavior of students.

Discussion

This study examined the effect of combined education of lectures and screening as animations on the knowledge, attitude, and behavior of elementary students regarding oral health in the City of Dehloran. The results revealed significant differences in the mean score of the students' knowledge, attitude, and behavior before and after the educational intervention. 36% of the subjects used dental floss. Karami [15], Ramezankhani [16], and Mazloomi [14], respectively, reported the dental floss uses of 51.7, 21, and 37.5 percent by the subjects.

Although statistically significant no relationship between gender and knowledge, attitude and behavior was found, the mean scores of attitude and behavior among the girls were higher than in the boys, and the mean scores of knowledge among the boys were higher than in the girls. In the study of Neda Babaei et al, girls had more knowledge than boys [17]. This could be due to the kind of intervention conducted by Babaei for students at the middle school level, because probably girls are more sensitive about their beauty including their oral health and beauty as compared to boys.

A number of studies have demonstrated the effect of teaching through various media on the oral health status. Mohammad-khah [11] showed that oral health education through

lectures and screenings increased the awareness, attitude, and practice of elementary students in Chabahar City, in which the effect of screening was found to be better than speech. In a study by Zarei [18], the mean scores of knowledge and practice to dental health increased after the intervention among the elementary school students in Qazvin Province. Babaei [17] also detected the impact of oral health education on raising the awareness of secondary school students in Hamadan. Karami [15] confirmed the effect of education based on a health belief model (HBM) on the changes in oral health behavior in the fifth grade students in Ahvaz City. Bourke studied Australian students and concluded that oral health training through theater had tremendous impacts on improving their attitude and performance [19]. An increase also occurred in the awareness of tenyear old children as a result of health education in a study by Worthington [20]. In a study by Athraa on128 students aged 11-12 years in Iraq, it was also observed that training using video and photo tutorials had an impact on the oral health of students [21], and these results were consistent with the results of the study by Hussain conducted on the students in the City of Erbil, and with the findings of Bankole in Nigeria [22,23].

Changes in oral health behaviors are related to visual technologies (photos and video clips)

and lead to reinforcing the impact of learning because seeing and hearing have high impact on behavior [24].

It seems that health education in accordance with new practices away from tensions based on childish exhilaration is a shortcut to achieve the goal of promoting children's oral health. In the present study, the relationship between the participants' classroom and their behavior was significant, in the sense that the fifth-grade students had better preventive behaviors. In this study, there was no significant relationship between the education level of parents and the knowledge, attitude and behavior of students. These results were consistent with the results of Babaei [17] and colleagues on middle school students in Babol.

The results of this study suggest that gathering students' comments on training practices at early ages, along with the use of attractive methods for educational content delivery such as video display, can promote their knowledge, attitude, and behavior concerning oral and dental health. The researchers were able to simultaneously compare the impacts of several training methods such as lectures, film screenings, or using methods such as painting and wall newspapers.

Conclusion

The positive results of educational intervention indicate that education through appropriate

methods can improve students' behavior in the field of oral health. Screening is effective in improving students' knowledge and skills as a training method. Therefore, it is suggested to have more emphasis on training oral hygiene during school ages.

Conflict of interest

The authors state that there is no conflict of interest.

Acknowledgements

The researchers deem necessary to thank sincerely the collaboration of all students participating in the project, as well as the managers and teachers at the study schools.

Funding/Support

This article is the result of HSR research project (No. 909405), which has been implemented by the support from Ilam University of Medical Sciences, Ilam. Iran.

References

- Mazloomi-Mahmoodabad SS, Taghavi AM, Barkhordari A, Alidousti F. Effect of role modeling through theater show in oral health education. J Islam Dent Assoc 2009; 21(2): 138-42. [In Persian]
- 2. Pakpour A, Heidarnia A, Hajizadeh E. Oral health status and its determinants in a sample of Iranian adolescents in Qazvin. J

Med Sci Qazvin Univ 2010; 15(3): 55-60. [In Persian]

- Fallahi A, Ahmadi F, Ghofranipour F, Malekafzali B, Hajizadeh E. Causes of dental caries from the perspectives of adolescents: A qualitative study. J Dent Sch 2013; 31(4): 224-33.
- Shariyat E, Mohammad Nejad E, Abotalebi G. Evaluation of Oral Health in School Students. JHC 2012; 13(3): 35-40. [In Persian]
- Nithila A, Bourgeois D, Burmes DE, Nurtomaa H. WHO global oral bank. 1986-96: An overview of oral health surveys at 12 years of age. Bulletin of World Health Organization 1998; 76: 237-44.
- Esm Hoseini G. Mouth and Teeth Health Care in Islam. UNMF 2005; 3(1): 24-34. [In Persian]
- Hobdell M, Peterson PE, Clarkson J, Johanson N. Global goals for health 2020. Int Dent J 2003; 53(5): 285-8.
- Keikhaee R, Rakhshani F, Izadi S, Hashemi Z. Survey of oral health behaviors and its associated factors in female students of primary schools in Zabol based on health belief model. J ZBMU 2013; 4(2): 33-41. [In Persian]
- Vahid Golpayegani M, Ansari G, Shirazi M. The knowledge, attitude and practice of grade 5 students regarding oral hygiene. J IAPD 2012; 6(13): 36-45. [In Persian]

- 10. Jabarifar SE, Eslamipoor F, Kavuani N, Mosavifar SA. Comparing the effect of rendering the film and lecture with charts and attitudes related to oral health knowledge and practice of secondary school students. J IAPD 2011; 5(11): 7-12. [In Persian]
- 11. Mohamadkhah F, Amin Shokravi F, Faghihzadeh S, Babaie Heydarabadi A, Kazembeigi F, Maghsoodi R. Comparison of Two Methods of Dental Health Education Lectures and Film Screenings on Knowledge, Attitude and Practice of Students. SJIMU 2012; 20(4): 67-75. [In Persian]
- Birang R, Shakerian K, Yazdanpanah Samani F, Nadimi M. The Effect of Education by Visual Media on Oral Health Promotion of Students. AMUJ 2007; 9(5): 1-6.
- 13. Baghiani Moghadam MH, Mazlomi S, Sharifirad GH. Health Education. 2nd Edition, Tehran: Asare Sobhan, 2009; p: 49.
- 14. Mazloomi Mahmoodabad SS, Roohani Tanekaboni N. Survey of some related factors to oral health in high school female students in Yazd, on the basis of health behavior model (HBM). JBUMS 2008; 15(3): 40-8. [In Persian]
- 15. Karami K, Shakerinejad G, Kabiry B. Effect of education based on health belief model on the alteration of oral health behaviors among students. Ilam University

of Medical Scinces 2014; 21(7): 134-41. [In Persian]

- 16. Ramezankhani A, Mazaheri M, Dehdari T, Movahedi M. Relationship between health belief model constructs and DMFT among five-grade boy students in the primary school in Dezfool. Scientific Med J 2011; 10(2): 221-28. [In Persian]
- 17. Babaee N, Kardan K, Aghazadeh F, Noribayat Sh. Effect of Oral and Dental Hygiene Education on the Knowledge of Caries Preventive Behaviors in the Guidance School Students. JBUMS 2012; 14(1): 83-7. [In Persian]
- 18. Zarei F. Effects of health education on oral health through role playing and painting on awareness and function of children. J Qazvin Univ Med Sci 2010; 14(1): 92-4.
- 19. Bourke LF. The use of theatre in dental health education. Aust Dent J 1991; 36(4): 301-10.
- 20. Worthington HV, Hill KB, Mooney J, Hamilton FA, Blinkhorn AS. A cluster randomized controlled trial of a dental health education program for 10-year-old children. J Public Health Dent 2001; 61(1): 22-7.
- 21. Mahmood AA. The effect of oral health educational pictures and video on periodontal health and behavior of school children. Iraqi Dental Journal 2016; 38(2): 63-9.

- 22. Hussain SR. Effectiveness of dental health education program on periodontal health status of nursery school children in Erbil City. Zanco J Med Sci 2012; 16(3): 175-9.
- 23. Bankole OO, Ibiyemi O, Oke GA. A dental health education video for Nigerian

children in the Yoruba Language. Afr J Biomed Res 2011; 14(1): 77-9.

24. Stina APN, Zamarioli CM, Carvalho EC. Effects of an educational video on the oral hygiene of patients with hematologic disorders. Rev Eletr Enf 2014; 16(2): 304-11.